

© EP0000 / EPO

PN-JP57065184 A 19820420

PD-1982-04-20

PR-JP19800141406 19801009

OPD-1980-10-09

TI-STABLE IMMOBILIZED ADENYLATE
KINASE COMPOSITE AND ITS
PREPARATION

IN-IMAHORI KAZUTOMO; NAGATA
KAZUHIKO; NAKASHIMA HIROSHI

PA-IMAHORI KAZUTOMO; RIKAGAKU
KENKYUSHO; UNITIKA LTD

IC-C12N9/12 ; C12N11/00

© WPI / DERWENT

TI-Heat resistant and stable adenylate
kinase - obtd. by cultivation of Bacillus
stearothermophilus

PR-JP19800141406

19801009;JP19800141405 19801009

PN-EP0050007 A 19820421 DW198217

Eng 056pp

-JP57065181 A 19820420 DW198221 000pp

-JP57065184 A 19820420 DW198221 000pp

-DK8104459 A 19820524 DW198224 000pp

-CA1173768 A 19840904 DW198440 000pp
-EP0050007 B 19860205 DW198606 Eng
000pp
-DE3173725G G 19860320 DW198613
000pp
-US4584272 A 19860422 DW198619 000pp
-JP1054995B B 19891121 DW198950
000pp
-JP2005391B B 19900201 DW199009
000pp
-DK165702B B 19930104 DW199306
C12N9/12 000pp
PA - (IMAH-I) IMAHORI K
-(RIKA) RIKAGAKU KENKYUSHO
-(NIRA) UNITIKA LTD
C-C12N9/12 C12R1/07
IC-C12N9/12 ;C12N11/02 ;C12P19/32
;C12R1/07
IN-IMAHORI K; NAGATA K; NAKAJIMA H;
HIROSHI N; KAZUHIKO N; KAZUTOMO I

AB - EP--50007 Heat-resistant **adenylate kinase** (AK) retaining at least 80% of its original activity after incubation in buffer soln. at 50 deg.C for 15 mins. is new.

- The AK is obtd. by cultivating a Bacillus strain, esp. B. stearothermophilus, in a nutrient medium. The cultivation is pref. effected continuously under conditions such that the dilution ratio is at least 0.9 of the max. specific growth rate (l/hr).

- The AK has higher and more prolonged stability than the **enzymes** obtd. previously, and it can be readily **immobilised** to give a prod. having good operational properties. The AK, esp. when **immobilised**, is useful in the prodn. of ATP from ADP or AMP in bioreactor systems.

EPAB - EP--50007 Heat-resistant **adenylate kinase** (AK) retaining at least 80% of its original activity after incubation in buffer soln. at 50 deg.C for 15 mins. is new.

-The AK is obtd. by cultivating a Bacillus strain, esp. B. stearothermophilus, in a nutrient medium. The cultivation is pref. effected continuously under conditions such that the dilution ratio is at least 0.9 of the max. specific growth rate (l/hr).

-The AK has higher and more prolonged stability than the enzymes obtd. previously, and it can be readily immobilised to give a prod. having good operational properties. The AK, esp. when immobilised, is useful in the prodn. of ATP from ADP or AMP in bioreactor systems. (56pp)

USAB-US4584272 Heat-resistant adenylate kinase is produced, by (a) cultivating a bacterium of the genus Bacillus; and (b) collecting prod. whose activity after incubation in a buffer soln. at 50 deg.C for 15 mins. is 80% or more of original activity before incubation.

- Pref. cultivation is performed continuously such that dilution rate D , is $0.9 (\mu)_{\max}$ or more, where $(\mu)_{\max}$ is the max. specific growth ratio (l/hr) of bacterium under continuous cultivation. Bacterium is *B. stearothermophilus*. Enzyme prod. is purified and/or immobilised on a water-insoluble carrier by a covalent bond.

- ADVANTAGE - Can be stored for a long period of time. (12pp)

OPD - 1980-10-09

CT - 5.Jnl.Ref; No-SR.Pub; 3.Jnl.Ref

DS - CH DE FR GB IT LI SE

AN - 1982-33470E [17]

PN - JP57065184 A 19820420

PD - 1982-04-20

AP - JP19800141406 19801009

IN - IMAHORI KAZUTOMO; others: 02

PA - IMAHORI KAZUTOMO; others: 02

TI - STABLE IMMOBILIZED ADENYLATE KINASE COMPOSITE AND ITS PREPARATION

AB - PURPOSE: Heat-resistant adenylate kinase is immobilized by binding or adsorbing it to a support to produce immobilized adenylate kinase with long-lasting stability as well as improved operability on immobilization and increased economical efficiency.

- CONSTITUTION: A strain in Bacillus stearothermophilus, whose optimal growing temperature is about 50-85 deg.C, is cultured and the cells produced are treated in a buffer solution at about 50 deg.C for about 15min to collect heat-resistant adenylate kinase of more than 80% activity. The adenylate kinase is bound to or adsorbed on a water-insoluble support to produce immobilized adenylate kinase composite.

SI-C12N9/12

I-C12N11/00